

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/259783691>

# Human Health Risks: Impact of Pesticide Application

Article · July 2013

---

CITATIONS  
10

READS  
210

1 author:



[Yasser El-Nahhal](#)

Islamic University of Gaza

68 PUBLICATIONS 1,328 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



The potential use of sewage sludge from sheikh ejleen TP for agriculture MSc thesis [View project](#)



Ecology [View project](#)

## Human Health Risks: Impact of Pesticide Application

Yasser El-Nahhal<sup>1</sup>, Abed Alkareem Radwan<sup>2</sup>

1. Dept of Earth and Environmental Sciences, Faculty of Sciences, The Islamic University-Gaza

2. Faculty of Nursing, The Islamic University-Gaza, Gaza, Palestinian National Authority (PNA), Palestine. Email: [y\\_el\\_nahhal@hotmail.com](mailto:y_el_nahhal@hotmail.com); Cell phone 00970599634708

### Abstract

This article reviews the application of pesticides in Gaza Strip, Palestine and discusses its associated health risks. This study is based on data collection and analysis.

Data showed that large quantities of pesticides are used in Gaza Strip and the quantities are increased annually. Analyzing the data indicates that large numbers of pesticides are used for controlling different types of pests. Some pesticides are restricted by law but are available in the local market. Classification of pesticides according to its biological activity indicates that insecticides are the largest uses among other pesticides. Reviewing the acute poisonous cases in health records indicates that the reported acute toxic cases were among local farmers in Gaza and the number of acute toxic cases increased annually indicating direct health risks associated with pesticide use. In addition, the increased number of congenital malformation among the newborns indicates indirect health risks. Moreover, the number of cancer cases in Khan Younis governorate indicates a positive association with pesticide use. Classification of pesticides according to WHO standards identified extreme toxic pesticides (e.g parathion), highly toxic (dichlorvos), moderately toxic (malathion) and less toxic ones. These Pesticides have a wide range of octanol-water partitioning coefficient ( $K_{ow}$ , log P) values (-0.8 - 6.6), which results in a variety of storage and transport patterns in human bodies. They may move from the storage sites (e.g fat bodies) via partitioning to other parts of the human body. A pesticide with high  $K_{ow}$  log P value (hydrophobic) such as Fenvalerate can be stored in fat containing particles and be released in milk secretion exposing fetus, mother, and infants to health risks. A satisfactory solution to these problems is the implementation of restriction measures and the performance of frequent pesticide residue analysis of food samples.

**Key words:** Contamination, health risk, toxicology, log  $K_{ow}$  value.

### 1. Introduction

Synthetic pesticides have been widely used for many years ago. For instance more than 18,000 pesticide products were licensed for use, and each year about 2 billion pounds of pesticides are applied to different places (Environmental Protection Agency 2002).

The application of pesticides has greatly contributed to the increase of world food and improved the human and animal health.

However, this application may have resulted in human health impacts. It has been documented that occupational exposure to pesticides resulted in an annual incidence of 18 cases of pesticide-related illness for every 100,000 workers in the United States (Calvert et al. 2004). Moreover the non- occupational exposure to pesticides may have side effects to non-target organism, including humans.

However, direct application of pesticide resulted in the accumulation of pesticides residues in vegetable and/or fruit samples in many countries. For instance, Malhat et al., (2013a) evaluated the residues of kresoxim methyl in apple and found that the terminal residues of kresoxim methyl were below the FAO/WHO maximum residue limit (MRL, 0.2 mg/kg) in apple when measured 14 days after the final application. In addition, Malhat et al. (2013b) determined etoxazole residues in in apples, strawberries and green beans by solid phase extraction clean-up and HPLC-DAD and found considerable concentrations but they were below Egyptian maximum residue limit. Mandour (2012) studied the contamination of ground water with insecticides in Egypt. His results confirmed the presence of high concentrations of insecticides, including organonitrogenous and organochlorine in tap drinking surface and ground water.

Osman et al., (2010) determined pesticide residues in vegetables marketed in Al-Qassim region, Saudi Arabia and found that 55% of the samples contained different residues in which 60% of the contaminated samples contained residues above the maximum residue limit,

Barakat et al., (2013) determined and identified 25 Organochlorine pesticides (OCPs) and 29 polychlorinated biphenyls (PCBs) in sediment samples collected from 34 locations in Lake Qarun. They concluded that according to the established sediment quality guidelines, OCPs (e.g.  $\gamma$ -HCH, endrin and chlordanes) would be more concerned OCP species for the ecotoxicological risk in Lake Qarun. Malhat and Nasr (2011) monitored the concentration of organophosphorus pesticides in fish samples from different tributaries of the Nile River in Egypt. They found considerable concentrations of Chlorpyrifos, cadusafos, diazinon, prothiphos and malathion were detected in fish tissues samples at level below the maximum residue limit.

Moreover, Mahmoud et al., (2013) determined organochlorine pesticides (OCPs) in the edible offal of Egyptian

buffalo and found that Tongues had the highest concentration of OCPs in a comparison to livers and kidneys in the examined samples. Somia and Madiha (2012) studied the pathological effects of dichlorvos and fenitrothion in mice and found that both insecticides caused degenerative changes in the liver and kidney of mice. Furthermore, Elserougy et al., (2013) detected organochlorine pesticides (OCPs) in placental and breast milk (BM) samples donated from healthy lactating mothers in Egypt. Amer et al., (2002) studied skin diseases among workers exposed to pesticides. Dermatological findings were positive in 78%, 76% and 54% of workers exposed to organophosphates, pyrethroids and carbamate pesticides respectively.

However, pesticides residues in food samples are hot topics in many countries. For instance in Egypt (Abbassy, 2001; Dogheim et al., 2001; Saleh et al., 1996), Jordan (Al-Nasir et al., 2001) Palestine (Safi et al., 2001a,b), Kuwait (Sawaya et al., 1999, 2000) Iran (Sdergern et al., 1978) and Pakistan (Tahir et al., 2001).

Furthermore, the indirect application of pesticides resulted in the accumulation of pesticide residues in many food samples. For instance: 1) honey samples (Boaxter and Saliba 1996, Al-Rifai and Akeel, 1997), 2) human milk (Dagher et al., 1999; Alawi et al., 1992; Saleh et al., 1998; Dogheim et al., 1996; Hashemy et al., 1977), 3) chicken eggs (Hashemy and Mosstofian 1979), and 4) fish (Paz 1976; Dogheim et al 1996a; El-Zorgani 1980).

Serious problems to non-target organisms and human beings leading to a number of pathological and disturbed biochemical processes (Abu-Murad 2000; Rashatwar and Ilyas 1984; Rambabu and Rao 1994; Soliman et al., 1997; Amr et al., 1997; Safi et al., 1993, Safi 2002) and toxicity to cyanobacteria (Kerkez 2012) and Fish (El-Najjar 2013).

The best-documented health effects involve the nervous system. The neurotoxic consequences of acute high-level pesticide exposure are well established: Exposure is associated with a range of symptoms as well as deficits in neurobehavioral performance and abnormalities in nerve function (Keifer and Mahurin 1997). Pesticide exposure may also be associated with increased risk of neurodegenerative disease, particularly Parkinson disease (Le Couteur et al. 1999).

This paper investigates the application of pesticides and surveys the acute and the non-acute toxic effects, and discusses their health-related risk.

## 2. Materials and Methods

### 2.1 Description of the study area

Gaza strip, the southern part of Palestine, a semiarid land of total area of 365 km<sup>2</sup> lies on the Mediterranean sea. About 181 km<sup>2</sup> is considered as a cropland and is dominated by vegetables (tomato, cucumber, pepper) and fruits (citrus, grapes). Farmers use intensive agriculture via green house technology. Due to the growing population, many of green houses are being established each year to provide sufficient amounts of vegetables.

These agricultural practices use large amounts of pesticides which may be harmful to the local population.

### 2.2 Data collection

**Pesticides data:** Several meetings with officials from Ministry of agriculture were conducted. In these meetings raw data of pesticides were handled. The data were classified according to historical information, target group, chemical class, and lethal dose that kill 50% of tested organisms (LD<sub>50</sub>).

**Pesticide markets in Gaza Strip** were surveyed by site visits to the shops. These shops were licensed by Ministry of agriculture in Gaza, Palestine.

Data on the acute and non-acute toxic effect of pesticides were collected from the health Department, local hospitals and with direct meeting with the technical staff working in the hospitals and/or private clinics. The data were screened, historically arranged and presented in a suitable way to provide a good data unit.

**2.3 Field visits:** several field visits were conducted to the agricultural land to view the application of pesticides by farmers. Some photos were collected.

## 3. Results and Discussion


### 3.1 Pesticide use in Gaza

The amounts of pesticides used in Gaza are shown in Table 1.

**Table 1.** Quantities (ton) of pesticides used in the past years

Year	Herbicide	Insecticide	Fungicide	Soil sterilants	Total
2002	9.86	95.58	123.91	303.00	532.35
2005	20.44	56.71	74.34	300.70	452.19
2006	24.94	55.27	55.65	111.60	247.46
2007	18.80	35.58	43.27	93.80	191.45
2008	18.20	49.65	42.20	193.60	303.65
2009	39.43	139.34	123.69	394.40	696.86
2010	18.78	144.68	99.63	162.40	425.49

It can be seen that the amounts of herbicides used in Gaza increased from year 2002 up 2006. Then sharp a reduction in herbicide use was observed in the following year except that in 2009. In contrast, the used amounts of insecticide and fungicide were high in the year 2002 and sharp reductions in the amount used were observed up to the year 2008 followed by a sharp increase in following year (Table 1). The amount of soil sterilants used followed the same trend. Furthermore, in Figure 1 the total amount of pesticide used in the last 10 years is presented.



**Figure 1.** Total amounts of pesticides used in the last ten years in Gaza

It is obvious from Figure 1 that the amount of soil sterilants is the largest among all, whereas the amount of herbicides is the lowest among all. The reason for that is the application rate of soil sterilants is several times larger than that of herbicide. Furthermore, it can be seen that the amounts of insecticides and fungicides are nearly equal to each other and several fold larger than that of herbicides. The explanation of these results is that farmers used herbicides one time a year whereas insecticides and fungicides are used several times a year due to intensive agriculture. Similar observations were seen in the surrounding countries (Mansour 2008). Furthermore, surveying the pesticide market provides detailed information on the current situation of pesticide names, and chemical groups and distribution of selling shops. More details are shown in Table 2-3.

**Table 2.** Distribution of pesticide shops in Gaza Strip

Place	selling shops
South zone (Rafah + Khan Younis)	53
Meddle zone	21
Gaza City	13
North zone	22

It is obvious that a large number of pesticide selling shops (109) are distributed all over Gaza Strip area. This is due to the agricultural nature of Gaza Strip. Beside the fact that large quantities of pesticides are used in the public health sector for controlling mosquitoes and flies by different municipalities in Gaza Strip. Table 3 provides more details on the active ingredients of pesticides

**Table 3.** Names of pesticide and their target groups used in Gaza.

Insecticides	Herbicides	Fungicides	Rodenticide	Soil Sterilants
Chlorfluazuron	Prometryne	Propineb	Bromadiolone	Methm Sodium
Abamectin	Bentazone	Penconazole	Coumatetralyl	Methy bromide
Methamidophos	Oxyfluorfen	Chlorothalonil	Metaldehyde	
Cyhexatin	2,4-D	Carbendazim		
Chlorpyrifos	Metamitron	Tetraconazole		
Novaluron	Oxadiazon	Sulphur		
Bifenazat	Pendimethalin	Trifloxystrobin		
*Fenamiphos	*Bromacil	Mancozeb		
*Imidacloprid	*Alachlor	Sulphur		
*Acetamiprid	*Atrazine	*Thiophanate methyl		
*Methyl-Parathion	*Terbutryne			
*Carbaryl	*Cyhexatin			

It can be seen that at least 12 insecticides, 12 herbicides, 10 fungicides, 3 rodenticide, and 2 soil sterilants are used annually in Gaza. However, reviewing the records of WHO and Ministry of Agriculture in Palestine, it appeared that some pesticides are internationally banned or have a restricted use but are still available for use in Gaza pesticide market. These pesticides are marked with a star (\*) to indicate a restriction on use. Furthermore, physicochemical properties and LD50 values of these pesticides are shown in Table 4. It is obvious that the solubility of the listed pesticides in Table 4 ranges from insoluble and/or few micrograms/l (e.g. Cypermethrin) to very soluble (solubility equal to 57.9 g/l, Methomoyl). This variety in solubility gives different leaching potentials and a possibility of groundwater contamination in Gaza. The solubility value of the pesticide above 30 mg/l may have the potential for groundwater contamination as previously reported (Williams et al, 1988). Accordingly, large fraction of the pesticides listed in Table 4 may contaminate the ground water in Gaza. However, there are other factors such as soil type  $K_{ow}$  log P. However, the value of  $K_{ow}$  log P ranges from (-0.8) to (6.6). These values in soil may result in a retention of the pesticides in the soil due to the possibility of binding to the soil organic matter and/or clay fraction. In the way around, the pesticide that has a high value of  $K_{ow}$  log P (e.g.6.6, Cypermethrin) may result in a variety of storage and transport patterns in human bodies and create health problems. Moreover, the value of Henry constant ( $K_H$ ) has a direct rule in the environmental fate of pesticides in Gaza. However,  $K_H$  values range from  $2 \times 10^{-10}$  (e.g Imidacloprid) to 3.5 as in Metaldehyde. These variations enabled different environmental fates of pesticides in Gaza. In addition the 50% degradation ( $DT_{50}$ ) ranges from few days 4d as in Matam Sodium to large number of days as in Atrazine (200d). This indicates persistence of some pesticides in Gaza. Moreover, the acute lethal dose that kill 50% of the tested organisms ( $LD_{50}$ ) of the listed pesticides in Table 4 ranged from extremely toxic pesticide (e.g parathion with  $LD_{50}$  equals to 2mg/kg), high toxic (Dichlorvos 50 mg/kg), moderate toxic (Cypermethrin with  $LD_{50}$  equals to 250 mg/kg) and less toxic one (Oxyfluorfen with  $LD_{50}$  equals to 5000 mg/kg). According to the values of  $LD_{50}$ , it is expected that some poisonous cases, direct effect of pesticide application, and congenital malformation cases occurred from one year to another. Table 5, provide cases of poisonous and congenital malformation. Studies of pesticide neurotoxicity have typically evaluated either the long-term effects of pesticide poisoning or the effects of occupational exposure. Pesticide poisoning may go undiagnosed, especially among farm workers with poor access to medical care (Moses et al. 1993) and particularly among women (London et al. 2002).

**Table 4.** Physicochemical properties of pesticides in Gaza Strip

Common name	Solubility in water	LD <sub>50</sub> mg/kg	K <sub>H</sub> Pa m <sup>3</sup> mol <sup>-1</sup>	K <sub>ow</sub> logP	DT <sub>50</sub>
Abamectin	10µg/l	10	2.7*10 <sup>-3</sup>	4.4	-
*Carbaryl	120 mg/l	850	-	1.59	28 d
Carbofuran	320 mg/l	8	-	1,52	30-60d
*Acetamiprid	4.25g/l	217	5.3*10 <sup>-8</sup>	0.88	15-30
Bifenazat	3.76 mg/l	5000	1*10 <sup>-3</sup>	3.4	80d
Chlorfluazuron	0.01mg/l	8500	5.41*10 <sup>-4</sup>	5.8	-
Cypermethrin	4µg/l	250	2*10 <sup>-2</sup>	6.6	
Chlorpyrifos	1.44	133	6.76*10 <sup>-1</sup>	4.7	
*Fenamiphos	400mg/l	10	9.2*10 <sup>-5</sup>		
Methamidophos	0.12 mg/l	200	1.6*10 <sup>-2</sup>	-0.8	
Azocyclin	0.12 mg/l	209	7*10 <sup>-2</sup>	5.3	
*Imidacloprid	0.61g/l	5000	2*10 <sup>-10</sup>	0.57	-
Diazinon	60 mg/l	1250	6.09*10 <sup>-2</sup>	3.3	
Malathion	145 mg/l	1375	-	2.75	
Parathion	11mg/l	2	0.0302	3.83	-
Dichlorvos	18g	50	2.58*10 <sup>-2</sup>	1.8	
Fenvalerate	10µg/l	451	-	5.01	
Oxyfluorfen	0.116mg/l	5000	-	4.47	292d
Methomoyl	57.9 g/l	30	2.13*10 <sup>-6</sup>	0.093	
Oxamyl	280 g/l	3.1	3.9*10 <sup>-8</sup>	-0.44	
Methidathion	200 mg/l	25	3.3 *10 <sup>-4</sup>	2.2	
Pendimethalin	0.3	1250	-	5.18	100 d
*Bromacil	807 mg/l	2000	-	1.88	150 d
*Alachlor	170.3mg/l	930	3.2 *10 <sup>-3</sup>	3.09	42-70d
*Atrazine	33mg/l	1869	1.5 *10 <sup>-4</sup>	2.5	105-200d
*Terbutryne	22mg/l	2500	1.5 *10 <sup>-3</sup>	3.65	70 d
*Cyhexatin	1 mg/l	540	-	-	-
Propineb	10mg/l	>5000	-	-0.26	-
Penconazole	730mg/l	2125	6.6 *10 <sup>-4</sup>	3.72	343 d
Chlorothalonil	0.81 mg/l	>5000	2.50 *10 <sup>-2</sup>	2.92	36 d
Carbendazim	29 mg/l	15000	3.6 *10 <sup>-3</sup>	1.38	>50 d
Tetraconazole	156 mg/l	1248	3.6 *10 <sup>-4</sup>	3.56	-
Trifloxystrobin	0.61 mg/l	>5000	2.3 *10 <sup>-3</sup>	4.5	8 d
Mancozeb	6.2 mg/l	>5000	-	-	15 d
Sulphur	Insoluble	>5000	-	-	-
*Thiophanate methyl	Insoluble	7500	-	1.5	28d
Bromadiolone	19 mg/l	1.125	5.55 *10 <sup>-5</sup>	-	-
Coumatetralyl	4 mg/l	16.5	1*10 <sup>-7</sup>	3.46	90 d
Metaldehyde	222 mg/l	283	3.5	0.12	-
Metam Sodium	722g/l	1800	-	<1	4d
Methy bromide	17.5g/l	*3.03	1.03 *10 <sup>-3</sup>	-	-

Source: Tomlin 2000.

Reviewing the health records showed that only the acute poisoning cases are admitted in the hospital for urgent medical care. These cases are reported in Table 5. The non-acute toxic cases are medically treated and sent back home. These cases are not registered in the health record. In addition, pesticide training courses on the safe and effective use of pesticides were conducted to the farmers. Accordingly, a reduction of poisonous cases among farmers is observed (Table 5). Moreover, the number of congenital malformation among children (Table 5) is increased annually. These data are in accordance with the increasing amount of pesticide used in Gaza (Table 1). These data are nearly similar to those reported in Egypt (Mansour 2008). The explanation of these results is that the application of pesticides resulted in accumulation of pesticides residues in vegetable and/or fruit samples (Malhat et al., 2013a,b, Osman et al., 2010) and contamination of groundwater (Mandour 2012). These residues have been shown to be stored in the human body and released again in milk secretion and placental samples in lactating mothers (Elserougy et al., 2013). It has been found that pesticide residues caused damage to livers and kidneys of Egyptian buffalo (Mahmoud et al., 2013), pathological degenerative changes in the liver and kidney of mice (Somia and Madiha, 2012) and skin diseases among workers exposed to pesticides (Amer et al., 2002). These data support our suggestion that the growing incidence of congenital malformation among the children is strongly correlated with pesticide application and/or consumption of food samples contaminated with pesticide residues.

**Table 5.** Number of poisonous and congenital malformation cases reported in Gaza Strip

Year	Acute poisonous	Congenital malformation
2008	49	20
2009	40	32
2010	32	45
2011	25	Na
Total	164	97

Na= not available

### 3.2 The dynamics of pesticides in the human body

The hydrophobic nature of pesticides (Table 4) allows them to accumulate in the adipose tissues and redistribute to different organs in the human bodies. The dynamics and the fate of pesticide in the human body are primarily controlled by a combination of factors such as transports, subsequent residence time, metabolism, concentration and physicochemical properties, sex, age, species, tissue and the integrity of the detoxifying organ especially liver and kidney. The most important physicochemical properties regarding the behavior of hydrophobic organic pesticides are aqueous solubility and octanol water partitioning coefficient ( $K_{ow}$ ).

The pesticides listed in Table 4 have a wide range of  $\log K_{ow}$  values starting from relatively hydrophilic, Oxamyl, ( $K_{ow}=-0.44$ ) to extremely hydrophobic, Cypermethrin, ( $K_{ow}=6.6$ ). Consequently, the pesticides of high  $K_{ow}$  values (e.g. Cypermethrin, Azocyclin) may be stored in human adipose tissues at different rates due to  $\log K_{ow}$  variations, which control differences in bio-transformation or excretion.

Due to the high fat content of the breast, it tends to be a suitable organ to accumulate and store highly hydrophobic organic contaminants. This may result in a very slow rate of elimination of pesticide (Matsumura, 1985). The partitioning of these chemicals in the adipose may protect them from metabolism by the active enzymes and make the reaction with DNA of the cells easier. The dynamics of pesticide in the human body may result in a distribution and/or transfer of pesticide from high concentration level (e.g. mothers) to a low level acceptor (e.g. stillborn, infants). Accordingly, the adipose tissue may act as a sink of the pesticide in the human body. An evidence of dynamic movements of contaminants is that their diffusion out through milk secretion. Previous analysis found high concentrations of chlorinated hydrocarbons (HCB, alpha, beta and gamma BHC, DDT, DDE) in milk secretions from nursing women (Saleh et al., 1996; Dagher et al., 1999; Saeed et al., 2000). Polychlorinated dibenzo-p-dioxines, polychlorinated dibenzo-furans and polychlorinated biphenyls were found in the human milk (Tunstra et al., 1994; Beck et al., 1994; Gonzalez et al., 1996; Kiviranta et al., 1999; Malisch et al., 2000; Yang et al., 2002). Further evidence of dynamic movements of pesticides (e.g. organochlorine compounds) is that their detection in the body of infants. For instance organochlorine were found in newborn in America (O'Leary et al., 1970), Canada (McLeod et al., 1971), Israel (Polishuk et al., 1977) India (Saxena et al., 1981, Siddiqui and Saxena 1985), Western Slovakia (Rosival et al., 1983) and Japan (Ando et al., 1985).

These findings suggest that the distribution and partitioning of pesticides from mother to infants may occur through milk secretion.

An alternative mechanism is that the pesticide moves from mother to stillborn through placenta. Previous reports found organochlorine contaminants in tissues of stillborns, and cord blood of the newborn babies and in the

human placenta (Curley et al., 1969; Polishuk et al., 1970). Polishuk et al. (1977) found greater concentrations of DDT, HCH and polychlorinated biphenyls in the lipids of human fetal blood than in the lipids of maternal blood. These findings reflect the prenatal exposure resulting from the transplacental passage of these contaminants. However, the movement of the contaminants through the placenta depends on its  $K_{ow}$  value. However, contaminants of low  $K_{ow}$  (2.31) such as carbaryl can cross the placenta and rapidly be distributed in all fetal tissues (Strother and Wheeler 1980).

However, the distribution of contaminants in the human body is controlled by  $K_{ow}$  value. Thus, the serum concentration of contaminants is positively correlated to the bulk concentration in the adipose tissues. This process may result in an even distribution of pesticide residues to different fat containing organs. Previous analysis (Petreas et al., 2004) did not report significant difference between the concentrations of a contaminant in the breast and in the abdominal adipose tissues. However, the fraction of the contaminants found in the blood serum may be adsorbed on the surface of the blood cells or be partitioned again with the lung tissue resulting in a leukemia, lung cancer, and other malignant tumors. Whereas the fraction of the contaminant remained in the adipose (e.g. breast) may develop a breast cancer.

However, the movements of the organic pesticide from the source (e.g. mother) to the acceptor (e.g. fetus, infant) may occur through one or more of the following modes:

1. Simple diffusion, pesticides of high hydrophobicity such as Chlorfluazuron (Table 4) can cross the placenta from regions of high concentrations (mother) to regions of low concentrations (fetus) up to equilibrium balance.
2. Facilitated diffusion, placental transfer or milk secretion, may also occur in accordance with the concentration gradients but more rapidly than simple diffusion.
3. Active transport, contaminant transfers opposite an electrochemical gradient and requires the expenditure of metabolic energy.
4. Special processes, these movements include, phagocytosis, pinocytosis and a break in the placental barriers. However, the movement of organic contaminants such as pesticides from mother to their fetus was early described by Ginsburg, (1971), Klaassen, (1986) and Salama, (1993) who reported similar mode of transfer.

### 3.3 Potential health risk associated with pesticide contamination

The dynamics of pesticide residues in the human body may be a matter of serious concern to public health especially for pregnant women, their fetus and their new babies. It has been shown that mother age can influence the accumulation of pesticide residues in circulating blood and its subsequent transfer to the fetus. A long life span may cause a great accumulation of contaminants (Salama et al., 1993). Mothers with high concentration of pesticide residues may transfer a fatal concentration to stillborn through cordial code which may result in a late abortions of pregnancy, or via milk recreation result in health risk to infants. These pesticide residues may have resulted in a serious health problems to non-target organisms and human beings leading to a number of pathological and disturbed biochemical processes (Abu Murad, 2000, Rashatwar and Ilyas 1984, Rambabu and Rao 1994, Soliman et al., 1997, Amer et al., 1997, Safi et al., 1993, Safi 2002). Olszyna-Marzys (1978) reported prenatal and postnatal intoxication of the infants with hexachlorobenzene in Turkey and methylmercury in Iraq due to the consumption of fungicide-treated seed wheat by pregnant and lactating mothers. The growing incidence of cancer cases among the children in Lebanon (average cancer cases 786 case/year) (El Saghier et al. 1998) might be attributed to the consumption of contaminated milk samples with pesticide residues (Elserougy et al., 2013). Furthermore, DDE and polychlorinated biphenyls (PCBs) were detected in blood serum samples collected from 43 females diagnosed with invasive adenocarcinoma of the breast, Port Said region of Egypt (Ahmed et al. 2002). The concentration of polychlorinated biphenyls in breast adipose tissue was positively correlated with the fish consumption in Canada (Paris-Pombo et al., 2003). Recent study (El-Nahhal and Harrarah 2013) showed an increased number of cancer cases in Khan Younis governorate. This indicates a positive association with pesticide use and/or agrochemicals due to the agricultural nature of Khan Younis governorate.

Extremely toxic pesticides, which have  $LD_{50}$  values less than 20 mg/kg, (Aldicarb, parathion, carbufuran, and monocrotophos) may be a cause of deaths to infants and children in Palestine (MOH 2001). A positive association was found between the pattern of food consumption and the increasing incidence of coronary heart disease in Saudi Arabia (Musaiger, 2002).

El-Hag et al. (2002) found that Carcinoma of breast, lymphomas and leukemias, in Saudi Arabia. However, the cases reported in Saudi Arabia may be attributed to the consumption of high fat food (Musaiger 2002) which might be contaminated with mycotoxin (Al-Julafi and Al-Fatih, 2001) and/or poly aromatic hydrocarbon (Al-Saleh and Al-Doush, 2002). These data are similar to those found in Palestine (Safi et al., 1993; Safi 2002) and correlated as consequences of overuse of banned pesticides.

## 4. Concluding remarks

Pesticides are commonly used in the agricultural sectors for pest control. Thus direct application of pesticides

may result in the contamination of food, soil and groundwater. The pesticides in Gaza are insecticides, herbicides, fungicides, rodenticides and soil sterilants. The quantities of insecticides and fungicides were higher than herbicides. Poisonous cases in Gaza are among the farm workers and the number of cases reduced annually. In contrast, the number of congenital malformation among the children is increased annually indicating indirect health risks.

The pesticides in Gaza belong to different chemical groups with a hydrophobic or hydrophilic nature. The hydrophobic ones may be accumulated in the fat tissue in the body and be built up to toxic or genotoxic levels. Pesticide can move in the human body by different ways and reach the fetus and be a cause of death. The movement of pesticide through milk secretion may be a cause of infant mortality due to the pre-maturity or low birth weight, malignant neoplasm for children or breast cancer. However, these contaminants may be of high risk exposure to women as shown by a number of breast cancer cases developed in Palestine (Safi 2002) and Egypt (Ahmed et al., 2002). However, the variation of physicochemical properties especially  $K_{ow}$  Log P values of pesticides (Table 4) make the partitioning and distribution quite different. Accordingly, the health risk is a pesticide dependent. It becomes obvious that a strong positive correlation may exist between the pesticide application and the growing incidence of non communicable diseases (cancer cases or heart disease) in Saudi Arabia, Egypt, Lebanon and Palestine.

The following steps may be implemented to reduce the health risk associated with pesticide application: 1) reduction of pesticide use to the minimum level; 2) pesticide residue analysis for vegetable, fruit samples before harvesting and marketing; 3) Analysis of the breast milk during the nursing period to avoid infant exposure to contaminated milk mainly with pesticide; and 3) Consumption of fat free food.

## 5. Acknowledgement

Dr. El-Nahhal acknowledges Alexander von Humboldt Stiftung/ Foundation Fellowship Grant no IV-PAL/1104842 STP, Germany

## Reference

- Amer M, Metwalli M, Abu el-Magd Y. Skin diseases and enzymatic antioxidant activity among workers exposed to pesticides. *East Mediterr Health J.* 2002;8(2-3):363-373.
- Abbassy, M. S. 2001. Pesticide residues in selected vegetables and fruits in Alexandria City, Egypt, 1997-1998. *Bull. Environ. Contam. Toxicol.* 67:225-232
- Abu Mourad, A. T. 2000. Hematological biomarkers of exposure in pesticide workers in Gaza Strip. Master thesis, Al-Quds University. Palestine.
- Ahmed, M. T., N. Loutfy and E. El Shiekh. 2002. Residue levels of DDE and PCBs in the blood serum of women in the Port Said region of Egypt. *J. Hazard. Mater.* 89:41-48
- Alawi, M. A., N. Ammari and Y. Al-Shuraiki. 1992. Organochlorine pesticide contamination in human milk samples from women living in Amman, Jordan. *Archi. Environ. Contam. Toxicol.* 23:235-239.
- Al-Julaifi, M. Z. and A. M. Al-Falih 2001 Detection of trichothecenes in animal feeds and foodstuffs during the years 1997 to 2000 in Saudi Arabia. *J. Food Prot.* 64:1603-1606.
- Al-Nasir, F., A. G. Jiries, M. I. Batarseh and F. Beese. 2001. Pesticides and trace metals residue in grape and home made wine in Jordan. *Environ. Monit. Assess.* 66:253-263.
- Al-Rifai, J, and N. Akeel. 1997. Determination of pesticide residues in imported and locally produced honey in Jordan. *J. Apic. Res.* 36:155-161.
- Al-Saleh, I. and I. Al-Doush. 2002. Gas chromatography-mass spectrometric determination of polycyclic aromatic hydrocarbons in five species of fish from three sites in the Arabian Gulf. *Int. J. Environ. Health Res.* 12 :193-200.
- Amr, M. M., Z. S. Halim and S. S. Moussa. 1997. Psychiatric disorders among Egyptian pesticide applicators and formulators. *Environ. Res.* 73:193-199.
- Ando, M.;Hirano, S.; Itoh, Y. 1985. Transfer of hexachlorobenzene (HCB) from mother to newborn baby through placenta and milk. *Arch. Toxicol* 56:195-201.
- Barakat AO, Khairy M, Aukaily I. Persistent organochlorine pesticide and PCB residues in surface sediments of Lake Qarun, a protected area of Egypt. *Chemosphere.* 2013 ;90(9):2467-2476.
- Beck H.; Drossa A.; Mather W. 1994. PCDD and PCDF exposure and levels in humans in Germany. *Environ. Health Perspect* 102: 173-185.
- Boxter, Y and R. Saliba. 1996. Determining residues of bromopropylate, coumpaphos and fluvalinate in lebanese honey after 12 years of treatment against varoatoses. *Proce. First Arab apicultural Congress Beirut Lebanon.*
- Calvert, G.M.; Plate D. K.; Das, R.; Rosales, R.; Shafey, O. & Thomsen, C. (2004). Acute occupational pesticide-related illness in the U.S. 1998-1999: Surveillance findings form the SENSOR-pesticides

- Program. *An. J. Ind. Med.*, 45., 14-23.
- Curley, A.; Copeland, M. F.; Kimbrough, R.D. 1969. Chlorinated hydrocarbon insecticides in organs of stillborn and blood of newborn babies. *Arch. Environ. Health*.19:628-632.
- Dagher, S. M., S. R. Talkouk, S. S. Nasrallah, I. R. Tannous and S. M. Mroueh. 1999. Relationship of dietary intake to DDE residues in breast milk of nursing mothers in Beirut. *Food Addit. Contam.* 16:307-312.
- Dogheim, S. M., A. S. Gad Alla and M. A. El-Marsafy. 2001. Monitoring of pesticide residues in Egyptian fruits and vegetables during 1996. *J. Assoc. Off. Anal. Chem.* 84: 519-531.
- Dogheim, S. M., A. S. Gad Alla, A. S. El-Syes, M. M. Almaz and E Y.Salama. 1996a. Organochlorine and organophosphorus pesticide residues in food from Egyptian local markets. *J. Assoc. Off. Anal. Chem.* 79:949-952.
- Dogheim, S.M., El-Z. Mohamed, A. S. Gad Alla, S. El-Saied, Y. S. Emel, M.A. Mohsen, M. S. Fahmy. 1996. Monitoring of pesticide residues in human milk, soil, water, and food samples collected from Kafr El-Zayat Governorate. *J. Assoc. Off. Anal. Chem.* 79:111-116
- El Hag, I. A., R. Katchabeswaran, C. L. Chiedozi and M. S. Kollur. 2002. Pattern and incidence of cancer in Northern Saudi Arabia. *Saudi Med. J.* 23:1210-1213.
- El-Nahhal, Y., Harrarah, S. (2013) Contamination of groundwater and associated disease: case study from Khan Younis Governorate, Gaza, PNA. *Journal of Environment and Earth Science* 3,5:147-154.
- EL-Najjar H. Sh. (2013) Eco-toxicity of Carbaryl, Chlorpyrifos, and Diuron, as Individual and in Mixtures on *Daphnia magna* and *Tilapia nilotica*. M.Sc Thesis, The Islamic University – Gaza – Palestine
- El Saghir, N. S., S. Adib, A. Mufarrij, S. Kahwaji, A. Taher, P. Issa and A. I. Shamseddine. 1998. Cancer in Lebanon: analysis of 10,220 cases from the American University of Beirut Medical Center. *J. Med. Liban.* 46:4-11.
- Elserougy S, Beshir S, Saad-Hussein A, Abouarab A. Organochlorine pesticide residues in biological compartments of healthy mothers. *Toxicol Ind Health.* 2013 ;29(5):441-448.
- El-Zorgani, G. A. 1980. Residues of organochlorine pesticides in fishes in Sudan. *J Environ. Sci. Health B.*15:1091-1098.
- Ginsburg, J. 1971. Placental drug transfer. *J. Ann. Rev. Pharmacol.* 11:387-391
- Gonzales M. J.; Jimenez B.; Hernandez L.M.; Gonnored M. F. 1996. Levels of PCDDs and PCDFs in human milk from population in Madrid and Paris. *Bull. Environ. Contam. Toxicol.* 56:196-204.
- Hashemy-Tonkabony, S. E. and F. Fateminassab. 1977. Chlorinated pesticide residues in milk of Iranian nursing mothers. *J. Dairy Sci.* 60:1858-1860.
- Hashemy-Tonkabony, S. E. and B.Mosstofian. 1979. Chlorinated pesticide residues in chicken egg. *Poult. Sci.* 58:1432-1434.
- Keifer M, Mahurin R. 1997. Chronic neurologic effects of pesticide overexposure. *Occup Med* 12:291–304.
- Kerkez S. (2012) Potential Toxicity of the Herbicides: Diuron, Diquat, Terbutryn and their Mixtures to Cyanobacterial Mats, Wadi Gaza, Palestine. M.Sc Thesis, The Islamic University – Gaza – Palestine
- Kiviranta H.; Purkunen R.; Vartiainen T. 1999. Levels and Trends of PCDD/Fs and PCBs in human milk in Finland. *Chemosphere* 38:311-323.
- Klaassen, C.D. 1986. Distribution, excretion, and absorption of toxicants. In: *The basic Science of Poisoning.* C.D. Klaassen, M.O. Amdur, J. Doull (eds). 3<sup>rd</sup> ed. Macmillan Publishing Co., New York, Toronto and London. p.33.
- Le Couteur DG, McLean AJ, Taylor MC, Woodham BL, Board PG. 1999. Pesticides and Parkinson's disease. *Biomed Pharmacother* 53:122–130.
- London L, De Grosbois S, Wesseling C, Kisting S, Rother H, Mergler D. 2002. Pesticide usage and health consequences for women in developing countries: out of sight, out of mind. *Int J Occup Environ Health* 8:46–59.
- Malisch R.; Bruns-Weller E.; Knoll A.; Furst P.; Mayer R.; Wiesmuller 2000. Results of “emergency quality control study” as confirmation of PCDD/PCDF-contamination of milk samples. *Chemosphere* 40:1033-1040.
- Mahmoud AF, Darwish WS, Morshdy AE, Eldaly EA, Ikenaka Y, Ishizuka M. (2013) Determination of organochlorine pesticides (OCPs) in the edible offal of Egyptian buffalo. *Jpn J Vet Res.*;61 Suppl:S58-63.
- Malhat F, Kamel E, Saber A, Hassan E, Youssef A, Almaz M, Hassan A, Fayz Ael-S. (2013a) Residues and dissipation of kresoxim methyl in apple under field condition. *Food Chem.*;140(1-2):371-4. doi: 10.1016/j.foodchem.2013.02.050. Epub 2013 Feb 24.
- Malhat F, Badawy H, Barakat D, Saber A. (2013b) Determination of etoxazole residues in fruits and vegetables by SPE clean-up and HPLC-DAD. *J Environ Sci Health B.* 2013;48(5):331-5. doi:

- 10.1080/03601234.2013.742371.
- Malhat F, Nasr I. Organophosphorus pesticides residues in fish samples from the River Nile tributaries in Egypt. *Bull Environ Contam Toxicol.* 2011;87(6):689-692.
- Mandour RA. Existence of insecticides in tap drinking surface and ground water in Dakahlyia Governorate, Egypt in 2011. *Int J Occup Environ Med.* 2012;3(1):27-32.
- Mansour S.A. 2008 Pesticide Impacts in Egypt. *In Reviews of Environmental Contamination and Toxicology*, D.M. Whitacre (ed.) *Vol 196*, 1 pp 1-51.
- Matsumura, F. 1985. Toxicology of insecticides. Plenum Press, New York and London. p.368.
- McLeod, H. A.; Grant, D. T.; Phillips, W. E. J. 1971. Pesticide residue and metabolites in placenta. *Can. J. Pub. Health.* 62:341-344.
- Ministry of Health (MOH), PCR and HMIS 2002. Ministry of Health, Ramalla, Cancer mortality, Cancer, 1995-2000. Palestine.
- Moses M, Johnson ES, Anger WK, Burse VW, Horstman SW, Jackson RJ, et al. 1993. Environmental equity and pesticide exposure. *Toxicol Ind Health* 9:913-959.
- Musaiger, A. O. 2002. Diet and prevention of coronary heart disease in the arab middle East countries. *Med. Princ. Pract.* 11:9-16.
- O'Leary, J.A., Davis, J. E.; Edmundson, W. F.; Reich, G. A. 1970. Transplacental passage of pesticides. *Am J.Obstet. Gynecol.* 107:65-69
- Olszyna-Marzys, A. E. 1978. Contaminants in human milk. *Acta Paediatr. Scand.* 67:571-576.
- Osman KA, Al-Humaid AM, Al-Rehiyani SM, Al-Redhaiman KN. Monitoring of pesticide residues in vegetables marketed in Al-Qassim region, Saudi Arabia. *Ecotoxicol Environ Saf.* 2010;73(6):1433-1439.
- Paris-Pombo A, Aronson KJ, Woolcott CG, King WD. 2003 Dietary predictors of concentrations of polychlorinated biphenyls in breast adipose tissue of women living in Ontario, Canada. *Arch Environ Health.* 58(1):48-54.
- Paz, J. D. 1976. Preliminary study of the occurrence and distribution of DDT residues in the Jordan watershed, 1971. *Pestic. Monit. J.* 10:96-100.
- Polishuk, Z. W.; Wassermann, M.; Wassermann, D.; Groner, Y.; Lazarovici, S.; Tomatis, L. 1970. Effects of pregnancy on the storage of organochlorine insecticides. *Arch. Environ. Health.* 20:215-217.
- Polishuk, Z. W.; Wassermann, D.; Wassermann, M.; Cucos, S.; Ron, M. 1977. Organochlorine compounds in mother and fetus during labor. *Environ. Research.* 13:278-284.
- Rambabu, P. J. and M. B. Roa. 1994. Effect of organochlorine and three organophosphate pesticides on glucose, glycogen, lipid and protine contents in tissues of freshwater snails *Bellamya Dissilis* (Muller). *Environ. Contam. Toxicol.* 53:142-148.
- Rashtwar, S. S. and I. Ilyas. 1984. Effect of phosphomidon in a freshwater teleost fish *Nemachelius denisonii* (Day) histopathological and biochemical studies. *J. Environ. Biol.* 5:1-8.
- Rosival, L., Szokolay, A., Grunt, J. 1983. Transplacental passage of pesticides into human embryo. *Czech. Med.* 6:1-6
- Saeed, T., N. W. Sawaya, N.Ahmad, S. Rajagopal, B. Dashti and A. Al-Awadhi.2000. Assessment of the levels of chlorinated pesticides in breast milk in Kuwait. *Food Addit. Contam.* 17: 1013-1018.
- Safi, J. 2002. Association between chronic exposure to pesticides and recorded cases of human malignancy in Gaza Governorates (1990-1999). *Sci. Total Environ.* 284: 75-84.
- Safi, J., N. Abu Foul, Y. El-Nahhal and A. El-Sebae. 2001a. Monitoring of pesticide residues on cucumber, tomatoes and strowberries in Gaza Governarates, Palestine. *Nahrung/Food* 46:34-49
- Safi, J., N. Abou Foul, Y. El-Nahhal and A. El-Sebae. 2001b. Monitoring of pesticide residues on green pepper, potatoes, Vicia faba, green bean and green peas in Gaza Governarate (PNA), Palestine. *J. Pest Cont. Environ. Sci.*9 (1):55-72.
- Safi, J., Y. El-Nahhal, A. S. Soliman and A. H. Elsebae. 1993. Mutagenic and carcinogenic pesticides used in the Gaza Strip Agricultural Environmental. *Sci. Total Environ.* 132: 371-380.
- Salama, K. A.; Bakry, M.N.; Abou-Donia, B.M. 1993. A review article on placental transfer of pesticides. *J. Occup. Med. Toxicol.* 2:383-397.
- Saleh, M. H., M. A. Afify and A. Kamel. 1998. Mother's milk protein profile, a possible biomarker for human exposure to insictecides. *J. Environ. Sci. Health B.* 33:645-655.
- Saleh, M. H., M. A. Afify, A. Ragab, G. El-Baroty, A. Kamel and A. K. El-Sebae. 1996. Breast milk as a biomarker for monitoring human exposure to environmental pollutants ACS. p 114-125. In: *Biomarkers for Agrochemicals and Toxic Substances: Application and Risk Assessment.* Washington D.C.
- Sawaya, W. N., A. F. Al-Awadhi, T. Saeed, A. Al-Omair, N. Ahmad, A. Husain, S. Khalafawi, H. Al-Omirah, B.

- Dashti, H. Al-Amiri and J. Al-Saqer. 1999. Kuwait's total diet study: dietary intake of organochlorine, carbamate, benzimidazole and phenylurea pesticide residues. *J. Assoc. Off. Anal. Chem.* 82:1458-1465.
- Sawaya, W. N., A. F. Al-Awadhi, T. Saeed, A. Al-Omair, A. , N. Ahmad, H. Al-Omirah, S. Al-Zenki, S. Khalafawi, J. Al-Otaibi and H. Al-Amiri. 2000. Dietary intake of organophosphate pesticides in Kuwait. *Food Chem.* 69: 331-338.
- Sdergren, A, R. Djirsarai, M. Gharbzadeh and A. Moinpour. 1978. Organochlorine residues in aquatic environments in Iran, 1974. *Pestic. Mont. J.* 12:81-86.
- Soliman, A. S., A. M. Smith, P. Cooper, K. Ismail, H. Khaled, S. Ismail, S. R. McPherson, A. I. Seifeldin and L. M. Bondy. 1997. Serum organochlorine pesticide levels in patients with colorectal cancer in Egypt. *Arch. Environ. Health.* 52:409-415.
- Saxena, M. C.; Siddiqui, M.K.J.; Bhargava, A.K.; Krishnamurti, C. R.; Kutty, D. 1981. Placental transfer of pesticides in humans. *Arch. Toxicol.* 48:127-134.
- Siddiqui, M.K.J.; Saxena, M. C. 1985. Placenta and milk as excretory routes of lipophilic pesticides in women. *Hum. Toxicol.* 4:249-255.
- Somia el-M, Madiha F. Pathological effects of dichlorvos and fenitrothion in mice. *Pathol Res Pract.* 2012 ;208(5):286-2891.
- Strother, A.; Wheeler, L. 1980. Excretion and deposition of <sup>14</sup>C-carbaryl in pregnant, nonpregnant, and fetal issues of the rat after acute administration. *Xenobiotica* 10:113-118.
- Tahir. S., T. Anwer, I. Ahmad, S. Aziz, A. Mohammad and K. Ahad. 2001. Determination of pesticide residues in fruits and vegetables in Islamabad market. *J. Environ. Biol.* 22:71-74.
- Tomlin, S. C. 2000. *The Pesticide Manual*. Twelfth Edition. British Crop Protection Council, Surry, UK.
- Tunstra L.G.M.T.; Tragg W.A.; Rhji J.A.V.; Spreng P.F.V.D 1994. The Dutch PCB/Dioxin study, development of a method for determination of dioxins, planar and other PCBs in human milk. *Chemosphere* 29:1959-1875.
- Yang J.; Shin D.; Park S.; Chang Y.; Kim D.; Ikonomou M. G. 2002. PCDDs, PCDFs, and PCBs concentrations in breast milk from 2 area in Korea: body burden of mothers and implications for feeding infants. *Chemospher* 46:419-428.
- Williams W., P. Holden, D. Parsons, M. Lorber. (1988) *Pesticides in Ground Water Data Base, Interim Report*. Washington: U.S. Environmental Protection Agency, Office of Pesticide Programs.
- U.S. EPA office of pesticides programs (2002). *F.Y 2002 Annual report* Willhington, DC: US.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

## CALL FOR PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

### IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

